

Date Sampled/Received: 15 Jan. 19 / 17 Jan. 19

A-1	Orga	nics -	Col	lorado
	~ 5		~ ~	or acc

Bob Yost 16350 WCR 76

Eaton

CO 80615

Product Identification	Compost
RR004 011519	

COMPOST TECHNICAL DATA SHEET

LABORATORY: Soil Control Lab	; 42 Hangar Way; Watsonville, CA 9507	6 <i>tel:</i> 831.724.5422	fax: 831.724.3188	
Compost Parameters	Reported as (units of measure)	Test Results	Test Results	
Plant Nutrients:	%, weight basis	Not reported	Not reported	
Moisture Content	%, wet weight basis	24.5		
Organic Matter Content	%, dry weight basis	30.7		
pH	units	8.72		
Soluble Salts (electrical conductivity EC ₅)	dS/m (mmhos/cm)	2.4		
Particle Size or Sieve Size	maxium aggregate size, inches	0.64		
Stability Indicator (respirometry	·)		Stability Rating:	
CO ₂ Evolution	mg CO ₂ -C/g OM/day	9.8	Un-Stable	
	mg CO ₂ -C/g TS/day	3.0	On-Stable	
Maturity Indicator (bioassay)				
Percent Emergence average % of control 93.3		93.3		
Relative Seedling Vigor	average % of control	84.9		
Select Pathogens	PASS/FAIL: per US EPA Class A standard, 40 CFR § 503.32(a)	Pass	Fecal coliform	
Trace Metals	PASS/FAIL: per US EPA Class A	Deve	As,Cd,Cr,Cu,Pb,Hg	
	standard, 40 CFR § 503.13, Tables 1 and 3.	Pass	Mo,Ni,Se,Zn	

Participants in the US Composting Council's Seal of Testing Assurance Program have shown the commitment to test their compost products on a prescribed basis and provide this data, along with compost end use instructions, as a means to better serve the needs of their compost customers.

Laboratory Group:	Jan19C	Laboratory Number: 9010456-1/1
Analyst: Assaf Sadeh	asy Sobel	www.compostlab.com



Date Sampled/Received: 15 Jan. 19 / 17 Jan. 19

	A-1	Organics	-	Col	lora	d
--	-----	-----------------	---	-----	------	---

Bob Yost 16350 WCR 76

Eaton

CO 80615

Product Identification	Compost
RR004 011519	

COMPOST TECHNICAL DATA SHEET

LABORATORY: Soil Control Lab	; 42 Hangar Way; Watsonville, CA 9507	6 <i>tel:</i> 831.724.5422	fax: 831.724.3188
Compost Parameters	Reported as (units of measure)	Test Results	Test Results
Plant Nutrients:	%, weight basis	%, wet weight basis	%, dry weight basis
Nitrogen	Total N	1.1	1.4
Phosphorus	P_2O_5	0.91	1.2
Potassium	K ₂ O	0.31	0.42
Calcium	Ca	1.2	1.5
Magnesium	Mg	0.20	0.27
Moisture Content	%, wet weight basis	24.5	
Organic Matter Content	%, dry weight basis	30.7	
рН	units	8.72	
Soluble Salts (electrical conductivity EC ₅)	dS/m (mmhos/cm)	2.4	
Particle Size or Sieve Size	% under 9.5 mm, dw basis	99.9	
Stability Indicator (respirometry	y)	Stability Rating:	
CO ₂ Evolution	mg CO ₂ -C/g OM/day	9.8 3.0 Un-Stable	
	mg CO ₂ -C/g TS/day		
Maturity Indicator (bioassay)			
Percent Emergence	average % of control	93.3	
Relative Seedling Vigor	average % of control	84.9	
Select Pathogens	PASS/FAIL: per US EPA Class A standard, 40 CFR § 503.32(a)	Pass	Fecal coliform
Trace Metals	PASS/FAIL: per US EPA Class A standard, 40 CFR § 503.13,	Pass	As,Cd,Cr,Cu,Pb,Hg
	Tables 1 and 3.		Mo,Ni,Se,Zn

Participants in the US Composting Council's Seal of Testing Assurance Program have shown the commitment to test their compost products on a prescribed basis and provide this data, along with compost end use instructions, as a means to better serve the needs of their compost customers.

Laboratory Group:	Jan19C	Laboratory Number:	9010456-1/1
Analyst: Assaf Sadeh	any Solel	www.compostlab.com	

SOIL CONTROL LAB

TEL: 831-724-5422 FAX: 831-724-3188 www.compostlab.com

Account #: 9010456-1/1-2355 Group: Jan19C #15 Reporting Date: February 4, 2019

A-1 Organics - Colorado 16350 WCR 76 Eaton, CO 80615 Attn: Bob Yost

Date Received: 17 Jan. 19
Sample Identification: RR004 011519
Sample ID #: 9010456 - 1/1

<u> </u>				
Nutrients	Dry wt.	As Rcvd.	units	Stability Indi
Total Nitrogen:	1.4	1.1	%	CO2 Evolution
Ammonia (NH ₄ -N):	1200	910	mg/kg	mg CO ₂ -C/g (
Nitrate (NO ₃ -N):	2.3	1.7	mg/kg	mg CO ₂ -C/g
Org. Nitrogen (OrgN):	1.3	0.98	%	Stability R
Phosphorus (as P ₂ O ₅):	1.2	0.92	%	
Phosphorus (P):	5300	4000	mg/kg	Maturity Indi
Potassium (as K ₂ O):	0.42	0.32	%	Compost:Ver
Potassium (K):	3500	2600	mg/kg	Emergence (
Calcium (Ca):	1.5	1.2	%	Seedling Vigo
Magnesium (Mg):	0.27	0.20	%	Descriptio
Sulfate (SO ₄ -S):	460	340	mg/kg	
Boron (Total B):	13	10	mg/kg	Pathogens
Moisture:	0	24.5	%	Fecal Coliforn
Sodium (Na):	0.13	0.096	%	Salmonella
Chloride (CI):	0.096	0.072	%	Date Tested: 17
pH Value:	NA	8.72	unit	
Bulk Density:	34	45	lb/cu ft	Physical Cor
Carbonates (CaCO ₃):	13	9.9	lb/ton	Total Plastic
Conductivity (EC5):	2.4	NA	mmhos/cm	Film Plastic
Organic Matter:	30.7	23.2	%	Glass
Organic Carbon:	17.0	13.0	%	Metal
Ash:	69.3	52.3	%	Sharps
C/N Ratio	12	12	ratio	Total
AgIndex	> 10	> 10	ratio	Total
M - 1 - 1 -	<u> </u>	EDA L''t		O: - D: - (- !)

	Stability Indicate	or:		
	CO2 Evolution		Respirometery	
	mg CO ₂ -C/g OM/day		9.8	
	mg CO ₂ -C/g TS/d	day	3.0	
	Stability Ratin	g	unstable	
	Moturity Indicat	- C	har Diagonay	
	Maturity Indicate		-	
	Compost:Vermic	uite (v:v)	1:2	
	Emergence (%)		93	
	Seedling Vigor (%	6)	85	
	Description of	Plants	healthy	
	Dothogono	Dogulto	Units	Dating
	Pathogens			Rating
	Fecal Coliform		MPN/g	pass
	Salmonella	> 3	MPN/4g	fail
	Date Tested: 17 Jar	. 19		
	Physical Contar	ninants**	% by weight	
	Total Plastic		< 0.1	
	Film Plastic		< 0.1	
'	Glass		< 0.1	
	Metal		< 0.1	
	Sharps		ND	
	Total		< 0.5	

Aginuex	<i>></i> 10	<i>></i> 10	Tallo
Metals	Dry wt.	EPA Limit	units
Aluminum (AI):	6600	-	mg/kg
Arsenic (As):	2.7	41	mg/kg
Cadmium (Cd):	< 1.0	39	mg/kg
Chromium (Cr):	17	-	mg/kg
Cobalt (Co)	2.5	-	mg/kg
Copper (Cu):	87	1500	mg/kg
Iron (Fe):	9900	-	mg/kg
Lead (Pb):	9.6	300	mg/kg
Manganese (Mn):	280	-	mg/kg
Mercury (Hg):	< 1.0	17	mg/kg
Molybdenum (Mo):	3.5	75	mg/kg
Nickel (Ni):	8.8	420	mg/kg
Selenium (Se):	1.2	100	mg/kg
Zinc (Zn):	210	2800	mg/kg

tion	
% by weight	
0.0	
0.0	
0.0	
0.1	
1.4	
6.0	
13.9	
78.6	
	0.0 0.0 0.0 0.1 1.4 6.0 13.9

*Greater than 4mm in size (Sharps greater than 2mm)

Analyst: Assaf Sadeh

*Sample was received and handled in accordance with TMECC procedures.

 Account No.:
 Date Received
 17 Jan. 19

 9010456 - 1/1 - 2355
 Sample i.d.
 RR004 011519

 Group:
 Jan19C No. 15
 Sample I.d. No.
 1/1
 9010456

INTERPRETATION: Page one of three

Is Your Compost Stable?

dry wt.

All Uses

Respiration Rate Biodegradation Rate of Your Pile						
9.8 mg CO2-C/	+++++++					
g OM/day		tely Unstable> <	Unstable	> < High For Mulch		
,				-		
Is Your Compost Mature?)					
is four compost mature?	-					
AmmoniaN/NitrateN ratio						
520 Ratio		+++++++++++++	++++++++++++	++++++++++++++++++++++++++++++		
	VeryMature> <	Mature		> < Immature		
Ammonia N ppm						
1200 mg/kg			+++++++++++	+++++++++++++++++++++++++++++++++++++++		
dry wt.	VeryMature> <	Mature		> < Immature		
Nitrate N ppm						
2.3 mg/kg	+					
dry wt.	< Immature > < Mature					
pH value						
8.72 units	+++++++++++++++++++++++++++++++++++++++	+++++++++++++				
Consumban Emanual	< Immature		> <	Mature > < Immature		
Cucumber Emergence 93.3 percent				+++++++++++++++++		
93.3 percent	< Immature	*************		> < Mature		
	· illillature			> Wature		
Is Your Compost Safe Re	garding Health?					
- 10 II'						
Fecal Coliform						
< 1000 MPN/g dry wt.	++++++ < Safe		ناك د ا	ah Facal Caliform		
Salmonella	< Sale		> < FILE	gh Fecal Coliform		
Greater than /4g dry wt.	++++++++++++++++					
Greater than 74g dry wt.	<safe (none="" detected)<="" td=""><td></td><td></td><td>Count(> 3 per 4 grams)</td></safe>			Count(> 3 per 4 grams)		
Metals US EPA 503	Coale (Horie detected)	7	riigir Gaimonciia (South(> 5 per 4 grains)		
Pass dry wt.	+++++++					
1 455 ary wa	<all metals="" pass<="" td=""><td>> -</td><td>< One or more Met</td><td>als Fail</td></all>	> -	< One or more Met	als Fail		
		•				
Does Your Compost Prov	<u>'ide Nutrients or Organic</u>	: Matter?				
Nutrients (N+P2O5+K2O)						
3.0 Percent	++++++++++++++++++	++				
dry wt.	<low> < Average</low>	9	> < High Nutrient	Content		
AgIndex (Nutrients / Sodium			P2O5+K2O) / (Na			
13 Ratio	++++++++++++++++++++++++++++++++++++++					
13 Ratio	Na & Cl > < Nutrient and Sodium and Chloride Provider > < Nutrient Provider					
Plant Available Nitrogen (Pa		ease for first season				
4 lbs/ton	+++++++++++					
wet wt.	Low Nitrogen Provider> <	Average Nitroge	en Provider	> <high nitrogen="" provider<="" td=""></high>		
C/N Ratio		- J		, <u>, , , , , , , , , , , , , , , , , , </u>		
12 Ratio	++++++++++++++++++	++				
	< Nitrogen Release > < N-	-Neutral > < N-Dem	and> < High Nitro	gen Demand		
Soluble Available Nutrients	& Salts (EC5 w/w dw)					
2.4 mmhos/cm	++++++++					
dry wt.	SloRelease> < Average Nutrient Release Rate > <high available="" nutrients<="" td=""></high>					
Lime Content (CaCO3)						
13 Lbs/ton	+++++++++++++++++++++++++++++++++++++++					
dry wt.	< Low > Averag	je >∣< High	Lime Content (as	CaCO3)		
What are the physical pro	perties of your compost	1?				
	,	<u></u>				
Percent Ash						
69.3 Percent	++++++++++++++++++++++++++++++++++++++					
dry wt.	< High Organic Matter	> < Average	> <	High Ash Content		
Sieve Size % > 6.3 MM (0.25						
1.5 Percent	+++++++++					

>|< Size May Restrict Uses for Potting mix and Golf Courses

Account No.: Date Received 17 Jan. 19 9010456 - 1/1 - 2355 Sample i.d. RR004 011519

Group: Jan19C No. 15 Sample I.d. No. 1/1 9010456

INTERPRETATION: Page two of three

Is Your Compost Stable? Respiration Rate

9.8 Moderate-selected use mg CO2-C/g OM/day

The respiration rate is a measurement of the biodegradation rate of the organic matter in the sample (as received). The respiration rate is determined by measuring the rate at which CO2 is released under optimized moisture and temperature conditions.

Is Your Compost Mature? AmmoniaN:NitrateN ratio

immature

520

Ammonia N ppm					
1200	immature				
Nitrate N ppm					
2.3	immature				
pH value					
8.72	immature				
·					

Composting to stabilize carbon can occur at such a rapid rate that sometimes phytotoxins remain in the compost and must be neutralized before using in high concentrations or in high-end uses. This step is called curing. Typically ammonia is in excess with the break-down of organic materials resulting in an increase in pH. This combination results in a loss of volatile ammonia (it smells). Once this toxic ammonia has been reduced and the pH drops, the microbes convert the ammonia to nitrates. A low ammonia + high nitrate score is indicative of a mature compost, however there are many exceptions. For example, a compost with a low pH (<7) will retain ammonia, while a compost with high lime content can lose ammonia before the organic fraction becomes stable. Composts must first be stable before curing indicators apply.

Cucumber Bioassay

93.3 Percent Cucumbers are chosen for this test because they are salt tolerant and very sensitive to ammonia and organic acid toxicity. Therefore, we can germinate seeds in high concentrations of compost to

measure phytotoxic effects without soluble salts being the limiting factor. Values above 80% for both percent emergence and vigor are indicative of a well-cured compost. Exceptions include very high salts that affect the cucumbers, excessive concentrations of nitrates and other nutrients that will be in range when formulated to make a growing media.

Is Your Compost Safe Regarding Health? Fecal Coliform

< 1000 / g dry wt. Fecal coliforms can survive in both aerobic and anaerobic conditions and is common in all initial compost piles. Most human pathogens occur from fecal matter and all fecal matter is loaded in fecal coliforms. Therefore fecal coliforms are used as an indicator to determine if the chosen method for pathogen reduction (heat for compost) has met the requirements of sufficient temperature, time and mixing. If the fecal coliforms are reduced to below 1000 per gram dry wt. it is assumed all others pathogens are eliminated. Potential problems are that fecal coliform can regrow during the curing phase or during shipping. This is because the conditions are now more favorable for growth than during the composting process.</p>

Salmonella Bacteria

Greater than 3 / 4g dry wt. Salmonella is not only another indicator organism but also a toxic microbe. It has been used in the case of biosolids industry to determine adequate pathogen reduction.

Metals

Pass The ten heavy metals listed in the EPA 503 regulations are chosen to determine if compost can be applied to ag land and handled without toxic effects. Most high concentrations of heavy metals are derived from woodwaste feedstock such as chrome-arsenic treated or lead painted demolition wood. Biosolids are rarely a problem.

Does Your Compost Provide Nutrients or Organic Matter? Nutrients (N+P2O5+K2O)

3.0 Average nutrient content

This value is the sum of the primary nutrients Nitrogen, Phosphorus and Potassium. Reported units are consistent with those found on fertilizer formulations. A sum greater than 5 is indicative of a compost with high nutrient content, and best used to supply nutrients to a receiving soil. A sum below 2 indicates low nutrient content, and is best-used to improve soil structure via the addition of organic matter. Most compost falls between 2 and 5.

Account No.: Date Received 17 Jan. 19 9010456 - 1/1 - 2355 Sample i.d. RR004 011519

Group: Jan19C No. 15 Sample I.d. No. 1/1 9010456

INTERPRETATION:

Page three of three

AgIndex (Nutrients/Na+CI)

High nutrient ratio Composts with low AgIndex values have high concentrations of sodium and/or chloride compared to nutrients. Repeated use of a compost with a low AgIndex (< 2) may result in sodium and/or chloride acting as the limiting factor compared to nutrients, governing application rates. These composts may be used on well-draining soils and/or with salt-tolerant plants. Additional nutrients form another source may be needed if the application rate is limited by sodium or chloride. If the AgIndex is above 10, nutrients optimal for plant growth will be available without concern of sodium and/or chloride toxicity. Composts with an AgIndex of above 10 are good for increasing nutrient levels for all soils. Most composts score between 2 and 10. Concentrations of nutrients, sodium, and chloride in the receiving soil should be considered when determining compost application rates. The AgIndex is a product of feedstock quality. Feedstock from dairy manure, marine waste, industrial wastes, and halophytic plants are likely to produce a finished compost with a low AgIndex.

Plant Available Nitrogen (lbs/ton)

4 Low N Provider Plant Available Nitrogen (PAN) is calculated by estimating the release rate of Nitrogen from the organic fraction of the compost. This estimate is based on the respiration rate, ammonia, and nitrate values. Despite the PAN value of the compost, additional sources of Nitrogen may be needed during he growing season to offset the Nitrogen demand of the microbes present in the compost. With ample nutrients these microbes can further breakdown organic matter in the compost and release bound Nitrogen. Nitrogen demand based on a high C/N ratio is not considered in the PAN calculation because additional Nitrogen should always be supplemented to the receiving soil when composts with a high C/N ratio are applied.

C/N Ratio

12 Indicates maturity As a guiding principal, a C/N ratio below 14 indicates maturity and above 14 indicates immaturity, however, there are many exceptions. Large woodchips (>6.3mm), bark, and redwood are slow to breakdown and therefore can result in a relatively stable product while the C/N ratio value is high. Additionally, some composts with chicken manure and/or green grass feedstocks can start with a C/N ratio below 15 and are very unstable. A C/N ratio below 10 supplies Nitrogen, while a ratio above 20 can deplete Nitrogen from the soil. The rate at which Nitrogen will be released or used by the microbes is indicated by the respiration rate. If the respiration rate is too high the transfer of Nitrogen will not be controlable.

Soluble Nutrients & Salts (EC5 w/w dw - mmhos/cm)

2.4 Average salts This value refers to all soluble ions including nutrients, sodium, chloride and some soluble organic compounds. The concentration of salts will change due to the release of salts from the organic matter as it degrades, volatilization of ammonia, decomposition of soluble organics, and conversion of molecular structure. High salts + high AgIndex is indicative of a compost high in readily available nutrients. The application rate of these composts should be limited by the optimum nutrient value based on soil analysis of the receiving soil. High Salts + low AgIndex is indicative of a compost low in nutrients with high concentrations of sodium and/or chloride. Limit the application rate according to the toxicity level of thesodium and/or chloride. Low salts indicates that the compost can be applied without risking salt toxicity, is likely a good source of organic matter, and that nutrients will release slowly over time.

Lime Content (lbs. per ton)

Average lime content Compost high in lime or carbonates are often those produced from chicken manure (layers) ash materials, and lime products. These are excellent products to use on a receiving soil where lime has been recommended by soil analysis to raise the pH. Composts with a high lime content should be closely considered for pH requirements when formulating potting mixes.

Physical Properties

Percent Ash

69.3 High ash content Ash is the non-organic fraction of a compost. Most composts contain approximately 50% ash (dry weight basis). Compost can be high in ash content for many reasons including: excess minerilzation(old compost), contamination with soil base material during turning, poor quality feedstock, and soil or mineral products added. Finding the source and reducing high ash content is often the fastest means to increasing nutrient quality of a compost.

Particle Size % > 6.3 MM (0.25")

1.5 May restrict use Large particles may restrict use for potting soils, golf course topdressings, seed-starter mixes, and where a fine size distribution is required. Composts with large particles can still be used as excellent additions to field soils, shrub mixes and mulches.

Appendix:			
Plant Available Nitrogen (PAN) calculations:		Estimated available nutrients for use when calculating application rates	
PAN = (X * (organic N)) + ((NH4-N) + (NO3-N))			lbs/ton (As Rcvd.)
X value =	If $RR < 2$ then $X = 0.1$		
	If RR = 2.1 to 5 then X = 0.2	Plant Available Nitrogen (PAN)	3.8
	If RR = 5.1 to 10 then X = 0.3	Ammonia (NH4-N)	1.82
If RR > 10 then $X = 0.4$		Nitrate (NO3-N)	0.00
Note: If C/N ratio > 15 additional N should be applied.		Available Phosphorus (P2O5*0.64)	11.6
RR = Respiration rate		Available Potassium (K2O)	6.3